

Turning sand into fertile soil with applied soil science



Clover Estate, SA

CLOVER ESTATE

Farm Facts

Mil Lel, 15km north east of Mount Gambier, SA South East

Enterprise: Cattle

Contract rearing of dairy heifers

Property Size: 100 hectares

Average Annual Rainfall: 700 mm

Elevation: 56 m

Motivation for Change

- ◆ Rising production costs and animal health concerns

Innovations

- ◆ Introducing biologically-based soil conditioners to balance the mineral and microbial status of the soil
- ◆ Strategic use of foliar fertilisers
- ◆ Ceasing chemical inputs
- ◆ Innovations commenced: 1995

Key Results

- ◆ Stock output increased by 33%
- ◆ 25% reduction in irrigation used per animal weight produced
- ◆ Infertile sand converted into fertile dark soils showing organic matter down to 60cm
- ◆ Thriving pastures with fewer weeds



David Clayfield has used applied soil science to create fertile, healthy soils from sand, in turn producing healthy pastures and healthy cows.

Faced with rising production costs and animal health concerns, David Clayfield made the link that improving soil health – physically, chemically and biologically – could address the cause of animal health and low productivity problems.

David eliminated the use of chemical and acid-based fertilisers, replacing them with biologically-based soil conditioners tailored to rectify deficiencies identified through soil tests.

Fifteen years on, yellow sands have turned into dark, healthy soils with a substantial increase in soil organic matter. Irrigation requirements have reduced from the region's typical seven to eight megalitres per hectare each year to five to six, subsequently reducing the energy used for pumping and distribution.

Pastures are thriving, animal health has improved and productivity has increased, while the veterinary bills have plummeted - positive results, all round.

Contact David Clayfield: dclayfield@bigpond.com



Clover Estate

David's grandfather came to Clover Estate in the mid 1930s to manage a large grazing estate. He bought a portion of it when the estate was subdivided for soldier settlement blocks after World War II. David's father grew up on the farm and has now retired, leaving David to run the property.

The typical soil in the region is five metres of sand overlying limestone. Native vegetation is low open forest dominated commonly by manna gum (*Eucalyptus viminalis*), brown stringybark (*Eucalyptus baxteri*) and blackwood (*Acacia melanoxylon*). Most of the native vegetation in south-east South Australia has been cleared to make way for farming and plantation forestry, but there is a small remnant left on the Clover Estate property.

There are no surface streams in the limestone plains area of south-east South Australia – rainfall infiltrates into the limestone, which forms an extensive shallow aquifer system. Dairying and cropping in the region depend on pumping irrigation water from this aquifer system.

David's grandfather and father operated Clover Estate as a dairy farm for many years, irrigating the pastures using water pumped from the shallow limestone aquifer underlying the property. After the dairy industry was deregulated the property proved to be too small to run viably as a milk producing dairy farm. Since 2000, the farm has been used to raise dairy heifers that are bred elsewhere in the district.

Heifers are reared on contract on a 'pay for weight gain' basis. The calves are brought to the property at four to five months age and within one year they have gained about 200 kg, i.e. a young dairy cow. The farm has the capacity to 'turn off' 600 to 700 heifers a year.

The current market for these young cows is the Republic of China. The heifers are shipped to China where they are mated upon arrival and begin producing milk before they are two years old.

A total of about 62 hectares of Clover Estate is irrigated and the rest is used for unirrigated grazing and cutting hay to provide feed during winter. A highly modified landscape, almost all native flora on Clover Estate has been replaced with introduced pasture species.



Clover Estate lucerne pastures which are interspersed with rye grass, clover and plaitain

Deficient Sandy Soils

South-east South Australia is made up of distinct land systems. The main underlying geological formation comprises limestone layers that have produced fertile soil, including the famous 'terra rossa' (red earth) soils of the Coonawarra wine region. Scattered in bands across the plains are low wide sand ridges, running roughly parallel to the present coastline. These sand ridges formed along ancient coastlines in the past few million years as the sea level rose and fell and the land surface was uplifted during ice ages. These sandy soils have low natural fertility.

When the large grazing properties were 'broken up' for farming and closer settlement, these sandy soil plots were not in demand. The sandy areas in private ownership were used for grazing, rather than cropping.

Clover Estate and many other properties in the area have been used for dairying, irrigated with groundwater pumped from the limestone aquifer, for many years. This groundwater is rich in calcium, which increases pH from natural levels of around 5, that is, slightly acidic, up to 7 to 8, which is slightly alkaline. This change causes nutrient imbalances and encourages growth of Yorkshire fog or velvet grass (*Holcus lanatus*), a perennial grass. *Holcus lanatus* is generally considered in Australia to be a weed of saline and waterlogging-prone sites and has little grazing value. The traditional way to redress the nutrient imbalances in the area is to use superphosphate and potash - potassium-rich fertiliser.

In 1992, a soil test showed organic carbon at 2.1%, total cation exchange capacity 3.5, and deficient levels of 17 macro and micro soil nutrients at Clover Estate. Prevailing advice was to continue to apply higher amounts of chemical fertiliser to maintain production. But high chemical and

fertiliser inputs and high water use were proving to be financially unsustainable. Furthermore, excessive weed competition had been developing despite regular use of knock-down and selective herbicides. Insecticides were used regularly for red-legged earth mite and other pests.

These problems, high levels of irrigation per animal production unit, on top of recurring animal health issues – mastitis, prolapse, sore feet – and concern about personal exposure to chemicals, led David to investigate more sustainable ways to improve soil and forage quality.

Identifying the Benefit of Healthy Soils

David had realised that management of animal health with medicines was only addressing the symptoms of animal health and performance, rather than the causes. Equally, chemical management was only creating recurring problems at extra cost. Conversely, improving soil health and mineral balance and availability in pasture would address the cause of animal health and low productivity problems.

David found that experimentation and mineral supplementation redressed some animal health problems. Success with this process indicated the strong link between animal nutrition and health. This prompted David to undertake further investigation to address animal health problems through the soil and fodder.

David's aim was to stop using chemical or acid-based fertilisers and pesticides. He wanted to improve soil fertility and water-holding capacity by adding biologically-based stimulants that increase soil organic matter.

He encountered challenges with adopting new methods, the initial impediment being a lack of information about alternatives to the 'single super' use way of farming. He was required to perform ongoing research, education and trial and error to identify his options and learn more about the links between soil and animal health.

David initially drew on information from Pat Coleby, a practitioner of animal health through soil health, which provided a useful guide. He also attended short courses in different approaches to fertiliser management, increasing his understanding and confidence in soil mineral and microbial management. David notes, "I have utilised interactions with agronomists and soil scientists from biological product supply companies to assist in fertiliser program management and planning".

David also read widely to supplement the courses and consultation. His collection of reference books include

Charles Walters and C.J. Fenzau's *Eco-farm*; Arden Andersen's *Science in Agriculture: Advanced Methods for Sustainable Farming*; and Pat Coleby's *Healthy Land for Healthy Cattle*.

An equally significant challenge was the lack of interest from other farmers, extending even to ridicule of some of the changes made and methods tried.

Regardless, David advises, "Work it out, stick to it, no matter what everyone thinks. With the experience and knowledge now available, you could build carbon and fertility in your soil in much less time than it took us to learn how by trial and error".

Turning Sand into Soil

Understanding the physical, chemical, and biological aspects of soil through applied soils science and putting this knowledge into practice over time, has been the fundamental innovative practice applied at Clover Estate.

"Testing the soil and observing the plants and animals, it became obvious that use of chemicals and acidic fertiliser were not a long-term solution", David says. He stopped using chemical fertilisers in the mid 1990s, five years before the switch from milking to raising heifers.

"The aim was to start with the soil health, by balancing the mineral and microbial status of soil, and the subsequent benefits in forage quality and animal health would follow."



Spraying biological liquid fertiliser



“The aim was to start with the soil health, by balancing the mineral and microbial status of soil, and the subsequent benefits in forage quality and animal health would follow.”

The process began with soil testing to assess the situation. Soil treatments were applied with the objective of addressing mineral balance, improve soil biological processes and overall soil fertility specific to paddock soil test results. The key ingredients are residue-digesting fungi and nitrogen-fixing bacteria, including strains of *Azotobacter*, *Bacillus*, *Pseudomonas*, and *Trichoderma*¹. Nutrients, carbohydrate and minerals are added to enable the bacteria and fungi to multiply in temperature-controlled 5000 litre aquaculture tanks. The minerals included a carbon, humic base to stimulate soil biological processes and organic matter building activity. The resulting brew is applied to the paddocks yearly at 50 litres a hectare.

In addition to soil building programs, a range of bio-fertilisers, such as foliar and fertigated (application through an irrigation system) mineral, microbial, kelp, fish, humic and fulvic fertilisers have been applied over the past 15 years. Leaf tests are taken to identify lacking nutrition, enabling application of nutrients to address imbalance in plants.

Animals’ performance and grazing preference, along with observing the plant species that are growing, are used as an additional guide as to excess and deficit minerals in the soil.

To target these treatments accurately, David has equipped himself with a refractometer, to measure the sugar concentration in plant sap, and other measurement devices and has soil tested by an agricultural laboratory in Adelaide.

In David’s words, the three rules for the soil remediation techniques he has developed and applied are “balance, balance and balance – carbon to nitrogen ratio, calcium to magnesium ratio, etc. ... look at the whole picture from soil to human health”.

The process began with soil testing to assess the situation. Soil treatments were applied with the objective of addressing mineral balance, improve soil biological processes and overall soil fertility specific to paddock soil test results. The key ingredients are residue-digesting




Above: Microbe brewing
Top: David Clayfield with bags of soil treatment used on the property

Healthy Soil Outcomes

The results are evident in the soil profiles of remnant and treated soils (see overleaf). Soil carbon has increased by 45% in the top 150mm, and potentially substantially more at depth.

Organic carbon measurements show an increase from 2.07% in 1992 to 2.92% in 2011.



David estimates that, since the mid 1990s, stock output has increased by 33% while he is using 25% less irrigation water per animal weight produced.

As is usual in the region, centre pivot systems are used at Clover Estate to irrigate with water pumped from the shallow limestone aquifer. Centre pivot irrigation uses sprays fitted on a boom that travels in a circle around the centre pivot point. Electric motors are used to drive the wheels that carry the boom. Two such systems operate at Clover Estate, one of about 300 metres radius, covering about 28 hectares, and the other a 330 metres radius, covering 34 hectares. A neutron probe is used to measure soil moisture levels and determine when watering is needed.

Due to the improved structure of the soils, David has found that his total water use is now around 5-6 MI/ha/year compared with the 7-8 MI/ha/year commonly used on sandy soils in the region. This is particularly noteworthy in relation to rainfall, with eight of the last ten years receiving below-average annual falls. As well as using less water, David's energy costs from pumping and driving the boom have halved.

Cessation of chemical weed control and David's soil management has ensured vigorous growth of preferred pasture species, combined with rotational grazing. The property is fenced into 50 paddocks, ranging from two to six hectares. The rotational grazing cycle also aims to break the life cycle of intestinal gut parasites.

David highlights, "Pasture plants grow better and there are now a lot fewer weeds. Lucerne varieties that had been abandoned due to poor performance by most typical, chemically managed properties perform exceptionally well on our property. Lucerne, rye grass, clovers, are the predominant species, along with a variety of herbs for balanced pasture. Animal health issues are primarily averted, due to the density and balance of the minerals in our pasture".

"Strategic use of selected foliar fertilisers... reduces or eliminates sap sucking insects attacking the pasture. We have found also that foliar fertilisers strengthen our desirable forage species, making them more competitive over weeds."

David believes that "Holistic management includes an appreciation of natural processes and an understanding of applied soil science. If soil biology functions to its best ability, the availability of minerals to plants will be at its best. If animals and people have access to adequate mineral diversity in diet, then disease is less ... and performance is better. Our farm enterprise is more profitable, and more enjoyable. We will leave an improved legacy."

"Our example has inspired other farmers, particularly dairy and graziers, to adopt aspects of soil management as we have. Looking at our soil profile, black to the depth that it is, speaks for itself. Most farmers appreciate that something is different and improved on our property. We are happy to share our journey, as it appears to give confidence to other food producers to take on similar management. Less chemicals affecting the environment, with improved quality of food for consumers is a better outcome than prior to adopting biological farming system."



Clover Estate centre pivot irrigation layout



FROM SAND TO SOIL

The typical condition of the original infertile sand is indicated by the profile under the patch of remnant native vegetation on the southern edge of the property, where no treatments have been applied (left). The results of 15 years of organic-based treatments are evident in the much darker colour, resulting from higher organic matter content down to 600mm where it used to be devoid of biological activity (right), which means that nutrient and water holding capacity are far higher.



¹ LawrieCo (2009), *Sustainable update winter 2009*, LawrieCo Sustainable Farming, Wingfield, South Australia.

This case study is an excerpt from the Soils for Life report:

Innovations for Regenerative Landscape Management: *Case studies of regenerative land management in practice*

REPORT SUMMARY

The Need for Change

Despite good practices of many of our land managers and farmers linked to some good science, the realities of an increasingly arid and degraded landscape will impact significantly not only on the productivity and viability of agricultural enterprises, but also on the health of our environment and the wellbeing of every Australian.

Landscape degradation is an issue of national and global concern. Landscape management practices including, but not limited to agriculture, forestry and fire have caused significant damage and in the process have altered the earth's natural biosystem. Consequently the precious resources of soil and water necessary to sustain life are being lost at unsustainable rates.

Unprecedented global challenges are arising in the face of this massive degradation of the landscape.

Soil erosion due to traditional agriculture is occurring at a rate between 10 and 100 times faster than the soil's natural formation process (pedogenesis)^{1,2}. Healthy soils are necessary to provide sufficient amounts of food with quality nutrition and fibre to meet global requirements.

Three billion people globally already have inadequate water and sanitation. It is assessed that 80% more water will need to be accessed by 2050 to feed the potential global population of more than nine billion³. Unless all limited soil and fresh water resources are understood and wisely managed, we are at risk of escalating social disruption and regional instability.

Even with its significant land area, Australia is not immune to the consequences of landscape degradation and increasing future needs. The realities of an increasingly arid and degraded landscape are already being experienced across the country. These include:

- increasing acidification, particularly in the south-east;
- declining soil health, caused by the loss of soil organic carbon (SOC);
- erosion;
- severe salinity;
- diminishing river flows;
- high evaporation and runoff rates;
- decreasing availability of groundwater; and
- reduced resilience to impacts of extreme and variable weather events such as drought, flood and fire.

The current state of the Australian natural landscape is further challenged by stresses from our changing climate, unsustainable management practices (such as reliance on high energy inputs), increased mining activity and urban expansion.

The national and global challenges being faced are interrelated and can be best met through a comprehensive coordinated approach focused on improved regenerative environmental management practices.

Landscape Regeneration for our Future

The key process drivers for landscape regeneration are **soil, water** and **vegetation**. Together in a natural system, supported by a constant flow of solar energy, these provide a regenerative cycle.

By restoring natural systems through improving landscape management practices, we can maximise water use efficiency, improve soil health, nutrient cycling and biodiversity of vegetation. A properly structured soil, with good levels of SOC, allows greater infiltration and retention of rainfall. Every gram of carbon in the soil can retain up to eight grams of water.

Currently, approximately 50% of rainfall on the Australian landscape is lost to evaporation due to poor soil structure and insufficient groundcover. By improving soil structure – particularly carbon – through increasing organic matter in the soil, we will be able to better capture and retain any rain that falls, making it available to plants for longer.

Through revegetation, groundcover is improved, and subsequently so is the quality of the soil, enhancing water infiltration. In turn, improved soil health and efficiency in water use contributes directly to the ability to support a biodiversity of vegetation and species.

If properly supported, this regenerative cycle can continue to sustain and improve the natural resource base and therefore landscape resilience and productivity.

Restoring these natural cycles and becoming more efficient in the use of natural resources is fundamental to the provision of sufficient food, fibre and water for a growing population. Business as usual is neither viable nor sustainable. Effective practical policies and actions are needed now.

Landscape Regeneration in Action

Innovative farmers are using high performance regenerative landscape management methods and fighting the trend of continued degradation of the landscape with its heavy reliance on external inputs. They are demonstrating sustainable, regenerative practices on their land. With relevant policies and incentives these practices could be extended successfully and quickly to involve a significant number of Australia's 135,000 farmers. Whilst there are always opportunities to learn more, enough is already known to take action now.



Soils for Life has documented some of these regenerative practices in 19 case studies across a range of locations and land-use types. Experiences shared by the 17 innovative farmers and two community organisations in the Soils for Life case studies demonstrate successful action being taken to restore the landscape. Due to the interrelated nature of soil, water and vegetation, benefits can be experienced across all process drivers regardless of the particular area of focus.

The Soils for Life case studies describe a range of techniques being used to obtain positive, regenerative outcomes, including:

- Applying organic composts, fertilisers and bio-amendments;
- Encouraging natural biological cycles and nutrient transfer;
- Implementing time-controlled planned grazing;
- Using grazing management and animal impact as farm and ecosystem development tools;
- Retaining stubble or performing biological stubble breakdown;
- Constructing interventions in the landscape or waterways to slow or capture the flow of water;
- Fencing off water ways and implementing water reticulation for stock;
- Investing in revegetation;
- Pasture cropping;
- Direct-drill cropping and pasture sowing;
- Changing crop rotations;
- Incorporating green manure or under-sowing of legumes;
- Managing for increasing species diversity;
- Controlling weeds through increased competition by desirable species;
- Reducing or ceasing synthetic chemical inputs; and
- Integrating enterprises.

PRINCIPLES FOR REGENERATIVE LANDSCAPE MANAGEMENT

Our case studies show that many different techniques can be applied to regenerate the landscape. Farmers and land managers commonly tailor a variety of methods to their own landscape and personal preferences. There is no single solution to landscape regeneration.

The following principles consistently emerge as underlying their regenerative practices – regardless of location or enterprise. These can be applied by other landholders as a basis for their own regeneration journey.

- Improve the structure of soil, through enhancing organic matter content
- Use and conserve rain where it falls
- Manage holistically
- Care about the land as a resource
- Commit to education and constant learning
- Search out communities of interest for help and advice
- Work on best land and extend from there
- Strive for maximum groundcover, for the majority of the time
- Manage times of plenty for times of shortage
- Reduce reliance on off-farm inputs
- Observe, measure and respond

Notes:

- 1 United Nations Environment Program, 2012, UNEP Year Book 2012: Emerging issues in our global environment, <http://www.unep.org/yearbook/2012>
- 2 Pimentel, D., 2006, 'Soil erosion: A food and environmental threat.' Environment Development and Sustainability, 8, pp119-137
- 3 Barlow, M., 2007, Blue Covenant: The Global Water Crisis and the Coming Battle for the Right to Water, McClelland & Stewart

CASE STUDY 6 - CLOVER ESTATE SA

Other case studies and the full *Soils for Life* report are available at: www.soilsforlife.org.au.

The production of the report was supported by funding provided by the Federal Government's *Caring for Our Country* initiative.

© Outcomes Australia, Soils for Life Program 2012

